

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® 647 Anti-Mouse CD1d Antibody[19G11]

Catalog Number: E-AB-F1032M

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

Isotype Rat IgG2b, κ **Clone No.** 19G11

Isotype Control Elab Fluor® 647 Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09842M]

Conjugation Elab Fluor® 647

Conjugation Information Elab Fluor[®] 647 is designed to be excited by the Red laser (627-640 nm) and detected

using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



C57BL/6 murine splenocytes are stained with Elab Fluor[®] 647 Anti-Mouse CD1d Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names Antigen-presenting glycoprotein CD1d1;CD1d.1;Cd1.1;Cd1d1

 Uniprot ID
 P11609

 Gene ID
 12479

For Research Use Only



Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Background

CD1d is a type I transmembrane protein and member of the MHC family, with a molecular weight ranging from 43-49 kD, depending on the glycosylation degree. CD1d is expressed by antigen presenting cells such as dendritic cells, monocytes, macrophages and B cells; also expressed by thymocytes and intestinal epithelial cells. CD1d present glycolipids to iNKT cells that recognize them by their Va14i TCR.