

## Recombinant Phospho-IKB alpha (Ser32, Ser36) Monoclonal Antibody

catalog number: **AN302083L**


**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human;Rat;Mouse
<b>Immunogen</b>	Recombinant human IKB alpha (phospho S32/36) fragment
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG, $\kappa$
<b>Clone</b>	A807
<b>Purification</b>	Protein A purified
<b>Buffer</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

### Applications Recommended Dilution

<b>WB</b>	1:500-1:2000
<b>IHC</b>	1:50-1:100



Immunohistochemistry of paraffin-embedded (A) Rat kidney, (B) Rat kidney, then the tissue was incubated with alkaline phosphatase (50U/ml) at 37°C for 1 hour using Phospho-IKB alpha (Ser32, Ser36) Monoclonal Antibody at dilution of 1:100.



Immunohistochemistry of paraffin-embedded (A) Mouse kidney, (B) Mouse kidney, then the tissue was incubated with alkaline phosphatase (50 U/ml) at 37°C for 1 hour using Phospho-IKB alpha (Ser32, Ser36) Monoclonal Antibody at dilution of 1:100.

### Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	Ice bag

### Background

NF-kappa-B inhibitor alpha (IKB alpha) inhibits the activity of dimeric NF-kappa-B/REL complexes by trapping REL dimers in the cytoplasm through masking of their nuclear localization signals. On cellular stimulation by immune and proinflammatory responses, IKB alpha becomes phosphorylated promoting itself ubiquitination and degradation, enabling the dimeric REL to translocate to the nucleus and activate transcription.

### For Research Use Only