

# Recombinant Human MAPK9 protein (His tag)

Catalog Number:PDEH100389



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	JNK-55;JNK2;JNK2A;JNK2ALPHA;JNK2B;JNK2BETA;p54a;p54aSAPK;PRKM9;SAPK;SAPK1a
<b>Species</b>	Human
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Phe 101-Arg 424
<b>Accession</b>	P45984
<b>Calculated Molecular Weight</b>	35.5 kDa
<b>Observed molecular weight</b>	41 kDa
<b>Tag</b>	N-His & C-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Mitogen-activated protein kinase 9 (MAPK9), also well known as c-Jun N-terminal kinase (JNK2), is a member of MAP kinase subfamily belonging to the protein kinase superfamily. The crystal structure of human JNK2 complexed with an indazole inhibitor by applying a high-throughput protein engineering and surface-site mutagenesis approach. A novel conformation of the activation loop is observed, which is not compatible with its phosphorylation by upstream kinases. This activation inhibitory conformation of JNK2 is stabilized by the MAP kinase insert that interacts with the activation loop in an induced-fit manner. It suggest that the MAP kinase insert of JNK2 plays a role in the regulation of JNK2 activation, possibly by interacting with intracellular binding partners. JNK2 deficiency leads to reduced c-Jun degradation,

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human MAPK9 protein (His tag)

Catalog Number:PDEH100389



thereby augmenting c-Jun levels and cellular proliferation, and suggests that JNK2 is a negative regulator of cellular proliferation in multiple cell types. JNK2 blocks the ubiquitination of tumor suppressor p53, and thus increases the stability of p53 in nonstressed cells. JNK2 negatively regulates antigen-specific CD8+ T cell expansion and effector function, and thus selectively blocking JNK2 in CD8+ T cells may potentially enhance anti-tumor immune response. Lack of JNK2 expression was associated with higher tumor aneuploidy and reduced DNA damage response. Additionally, the JNK2 protein could be a novel therapeutic target in dry eye disease, and may provide a novel target for prevention of vascular disease and atherosclerosis.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017