Recombinant Rat CCI2 protein (His Tag)

Catalog Number: PDMR100017



| Description | | | |
|----------------|--|--|--|
| Species | Rat | | |
| Mol_Mass | 13.6 kDa | | |
| Accession | P14844 | | |
| Bio-activity | Not validated for activity | | |
| Properties | | | |
| Purity | > 95% as determined by reducing SDS-PAGE. | | |
| Endotoxin | < 1.0 EU/mg of the protein as determined by the LAL method | | |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 | | |
| | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of | | |
| | reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. | | |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. | | |
| Formulation | Lyophilized from a 0.2 μ m filtered solution in PBS with 5% Trehalose and 5% | | |
| | Mannitol. | | |
| Reconstitution | It is recommended that sterile water be added to the vial to prepare a stock solution of | | |
| | 0.5 mg/mL. Concentration is measured by UV-Vis. | | |
| Data | | | |

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| KDa | М | R |
|-----|---|--------|
| 80 | - | - |
| 60 | | - 22 |
| | | |
| 40 | | - 6000 |
| | | |
| 30 | - | - 10 |
| | | |
| 20 | | |
| | | |
| 12 | | |

> 95 % as determined by reducing SDS-PAGE.

Background

The chemokine (C-C motif) ligand 2 (CCL2), also known as monocyte chemoattractant protein (MCP)-1 and small inducible cytokine A2 (SCYA2)), is a small cytokine that belongs to the CC chemokine family responsible for monocyte attraction. Its cognate receptor, CCR2, play a critical role in regulating nociceptive processes during neuropathic pain. Both CCL2 and CCR2 are implicated in induction of autoimmunity. CCL2 recruits monocytes, memory T cells, and dendritic cells to the sites of inflammation produced by either tissue injury or infection. Recently research also showed that CCL2 might be useful as a biomarker of fibrosis as well as a target for therapeutic intervention.

For Research Use Only