

# Phospho-MLKL (Ser358) Monoclonal Antibody

catalog number: E-AB-21338

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description


<b>Reactivity</b>	Human
<b>Immunogen</b>	Synthetic Peptide of Phospho-MLKL (Ser358)
<b>Host</b>	Mouse
<b>Isotype</b>	IgG
<b>Clone</b>	6E3
<b>Purification</b>	Protein A purification
<b>Conjugation</b>	Unconjugated
<b>buffer</b>	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein protectant and 50% glycerol.

## Applications


## Recommended Dilution

<b>IHC</b>	1:100-200
------------	-----------

## Data



Immunohistochemistry of paraffin-embedded Human colon carcinoma tissue with Phospho-MLKL (Ser358) Monoclonal Antibody at dilution of 1:200



Immunohistochemistry of paraffin-embedded Human skin tissue with Phospho-MLKL (Ser358) Monoclonal Antibody at dilution of 1:200

## Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

## Background

This gene belongs to the protein kinase superfamily. The encoded protein contains a protein kinase-like domain; however, is thought to be inactive because it lacks several residues required for activity. This protein plays a critical role in tumor necrosis factor (TNF)-induced necroptosis, a programmed cell death process, via interaction with receptor-interacting protein 3 (RIP3), which is a key signaling molecule in necroptosis pathway. Inhibitor studies and knockdown of this gene inhibited TNF-induced necrosis. High levels of this protein and RIP3 are associated with inflammatory bowel disease in children. Alternatively spliced transcript variants have been described for this gene. MLKL (Mixed Lineage Kinase Domain Like Pseudokinase) is a Protein Coding gene. Among its related pathways are Apoptosis and Autophagy and CDK-mediated phosphorylation and removal of Cdc6. GO annotations related to this gene include transferase activity, transferring phosphorus-containing groups and protein tyrosine kinase activity.

## For Research Use Only