TCEB2 Polyclonal Antibody

catalog number: E-AB-18260




Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | | |
|--------------|--|--|
| Reactivity | Human;Mouse;Rat | |
| Immunogen | Full length fusion protein | |
| Host | Rabbit | |
| Is otype | IgG | |
| Purification | Antigen affinity purification | |
| Conjugation | Unconjugated | |
| buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. | |

| Applications | Recommended Dilution |
|--------------|-----------------------------|
| IHC | 1:30-1:150 |

Data



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using TCEB2 Polyclonal Antibody at dilution of 1:40(×200)

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack,upon receipt,store it immediately at the |
| | temperature recommended. |

Background

This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on chromosomes 11 and 13.

For Research Use Only