Purified Anti-Mouse CD90 Antibody[M5/49.4.1], Functional Grade

catalog number: **E-AB-F12830**


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description


Reactivity	Mouse
Immunogen	Recombinant Mouse CD90 protein
Host	Rat
Isotype	Rat IgG2a, κ
Clone	M5/49.4.1
Purification	> 98%, Protein A/G purified, ≥ 95% monomer by analytical SEC
Endotoxin	< 1 EU/mg determined by LAL assay
Conjugation	Unconjugated
Buffer	Sterile PBS, pH 7.2. Contains no stabilizers or preservatives

Applications	Recommended Dilution
FCM	≤ 0.2 μg per million cells in 100 μL volume
Depletion	Reported in the literature

Data



C57/BL6 Mouse splenocytes were stained with 0.2 μg Purified Anti-Mouse CD90 Antibody[M5/49.4.1], Functional Grade (Right) and 0.2 μg Rat IgG2a, κ Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Rat IgG Secondary Antibody.



Monomer purity ≥95% as determined by analytical size-exclusion chromatography (SEC)

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze/thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.
Shipping	Ice bag

Background

CD90, also known as Thy-1, is a 28-30 kD GPI-linked membrane glycoprotein. It is a member of the immunoglobulin superfamily and has been shown to interact with CD45 in signal transduction during lymphocyte proliferation and differentiation. CD90 is expressed on hematopoietic stem cells, neurons, thymocytes, peripheral T cells, fibroblasts, stromal cells.

None (Azide-Free, Low Endotoxin) are perfectly suited to be used in culture or in vivo (for nonhuman studies) for functional assays blocking, neutralizing, activation or depletion where the presence of azide may damage cells or exogenous endotoxin may signal or activate cells.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com

Application References

Xiufang Weng, et al. Eur J Immunol. 2014 Dec;44(12):3646-57.