A Reliable Research Partner in Life Science and Medicine

Recombinant Casein Kinase 1 gamma 2 Monoclonal Antibody

catalog number: AN301832L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human;Rat;Mouse

Immunogen Recombinant human Casein Kinase 1 gamma 2 fragment

HostRabbitIsotypeIgG, κCloneA544

Purification Protein A purified


Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution

WB 1:1000-1:5000

IF 1:50 **FCM** 1:50-1:100


Data




Western Blot with Casein Kinase 1 gamma 2 Monoclonal Antibody at dilution of 1:5000. Lane 1: HeLa, Lane 2: 4T1, Lane 3: RAW 264.7, Lane 4: Rat brain

Immunofluorescent analysis of (4% Paraformaldehyde) fixed Hela cells using anti-Casein Kinase 1 gamma 2 Monoclonal Antibody at dilution of 1:50.

Observed-MW:50 kDa Calculated-MW:50 kDa





Immunofluorescent analysis of (4% Paraformaldehyde) fixed Flow cytometric analysis of mouse Casein Kinase 1 gamma
A549 cells using anti-Casein Kinase 1 gamma 2 Monoclonal 2 expression on Hela cells. Cells were stained with purified anti-Mouse Casein Kinase 1 gamma 2, then a Alexa Fluor

2 expression on Hela cells. Cells were stained with purified anti-Mouse Casein Kinase 1 gamma 2, then a Alexa Fluor 488-conjugated second step antibody. The histogram were derived from events with the forward and side light-scatter characteristics of intact cells.

Rev. V1.1


 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



Elabscience®



Flow cytometric analysis of mouse Casein Kinase 1 gamma 2 expression on A549 cells. Cells were stained with purified anti-Mouse Casein Kinase 1 gamma 2, then a Alexa Fluor 488-conjugated second step antibody. The histogram were derived from events with the forward and side light-scatter characteristics of intact cells.

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

The casein kinases (CK) belong to the serine/threonine kinases and can be subdivided further into either casein kinase 1 (CK1) or casein kinase 2 (CK2) families due to their high homology in their catalytic domains. In vertebrates, seven CK1 isoforms (α , β , γ 1, γ 2, γ 3, δ and ϵ) and several splice variants for CK1 α , δ , ϵ and γ 3 have been identified. They differ in length and sequence of the N-terminal (9–76 amino acids) and especially the C-terminal (24–200 amino acids) non-catalytic domain.

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.1