

## Purified Anti-Human CD56 Antibody[MY31]

catalog number: E-AB-F1270A

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

### Description


<b>Reactivity</b>	Human
<b>Immunogen</b>	Recombinant Human CD56 protein
<b>Host</b>	Mouse
<b>Isotype</b>	Mouse IgG1, $\kappa$
<b>Clone</b>	MY31
<b>Purification</b>	>98%, Protein A/G purified
<b>Buffer</b>	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

### Applications

### Recommended Dilution

**FCM** 2  $\mu$ g/mL (0.5 $\times$ 10<sup>6</sup>-1 $\times$ 10<sup>6</sup> cells)

### Data



Human peripheral blood lymphocytes cell were stained with 0.2  $\mu$ g Purified Anti-Human CD56 Antibody[MY31] (Right) and 0.2  $\mu$ g Mouse IgG1,  $\kappa$  Isotype Control (Left),

followed by Elab Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD16 PE-conjugated Monoclonal Antibody.

### Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	Ice bag

### Background

This gene encodes a cell adhesion protein which is a member of the immunoglobulin superfamily. The encoded protein is involved in cell-to-cell interactions as well as cell-matrix interactions during development and differentiation. The encoded protein has been shown to be involved in development of the nervous system, and for cells involved in the expansion of T cells and dendritic cells which play an important role in immune surveillance. Alternative splicing results in multiple transcript variants.

### For Research Use Only