

Ham's F-12K, powder

Cat. No. : PM150910P

Size : 5×1L / 1×10L / 1×50L / 100L / 500L

General Information

| | |
|--------------------|-----------------------|
| Product Form | Powder |
| D-Glucose | 1260 mg/L |
| Concentration | 11.09 g/L |
| HEPES | Negative [®] |
| L-Glutamine | 2 mM |
| NaHCO ₃ | Negative |
| Phenol red | 3 mg/L |
| Sodium pyruvate | 2 mM |
| Storage | 2-8°C, Shading Light |
| Shipping | Room Temperature |
| Expiration date | 36 months |

Background

The Ham's F-12K medium increases the concentration of amino acids and pyruvic acid on the basis of Ham's F-12, reduces the concentration of glucose, and modifies the composition and content of the salt.

The Ham's F-12K medium was initially designed to culture primary human hepatocytes and differentiated cells of rats and chickens.

The Ham's F-12K medium contains many kinds of amino acids, vitamins, inorganic salts and other ingredients for cell culture, but does not contain protein, lipids or any growth factors. Therefore, the product should be used with serum or serum-free additives.

Preparation method

1. The preparation water should be purified water, ultra-pure water or water for injection (WFI), and the water temperature should be controlled between 20-30°C during the preparation process.
2. Measure 90% of the final volume preparation water to the solution preparation system. Start stirring, and avoid generating bubbles. For example, if 1 L is required, add 900 mL of preparation water here. And it's recommended that the power output per unit volume (P/V) of the mixing system is greater than 10 W/m³.
3. Weigh the appropriate amount of powder according to the concentration of 11.09 g/L accurately, and add it to the container prepared in step 2. Stir for more than 20 minutes dissolve all powder completely.
4. After the solution is clear, add NaHCO₃ at a concentration of 2.5 g/L, continue stirring for 5-10 minutes until dissolved, then add ultra pure water to adjust the volume to the 100% of required.
5. If necessary, adjust the pH to 7.20-7.30 with 1 mol/L NaOH solution or 1 mol/L HCl solution. Since filtration will slightly increase the pH, the pH value here is lower than the target pH value (7.20-7.40).

6. The prepared solution should be sterilized using a 0.2 µm pore size filter membrane under positive pressure (ensure aseptic technique).
7. After filtration, a small amount of liquid culture medium can be taken for quality inspection, and use only after passing the test.
8. The filtered liquid medium should be used immediately or stored in glass bottles, culture medium bottles (PET), or single-use storage bags with an oxygen-barrier coating at 2-8°C away from light. The liquid medium has a shelf life of 1 year under these conditions.

Notes

1. Please wear a lab coat and use disposable gloves and a mask during operation.
2. To ensure the optimal performance of this product, please strictly adhere to the recommended storage conditions for its preservation.
3. This product is intended for scientific research exclusively or as a raw material in the production process, and must not be applied for clinical diagnosis or treatment.