

Purified Anti-Mouse IL-10 Antibody[JES5-16E3]

catalog number: E-AB-F1197A

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Mouse
Immunogen	Recombinant Mouse IL-10 protein
Host	Rat
Isotype	Rat IgG2b, κ
Clone	JES5-16E3
Purification	>98%, Protein A/G purified
Buffer	PBS, pH 7.2. Contains 0.05% proclin 300.

Applications Recommended Dilution

FCM	2 μ g/mL (1×10^5 - 5×10^5 cells)
------------	---

Data



HEK293T cells transfected with pcDNA3.1 plasmid encoding Mouse IL-10 gene were stained with 0.2 μ g Purified Anti-Mouse IL-10 Antibody[JES5-16E3] (Right) and 0.2 μ g Rat IgG2b, κ Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Rat IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

Major immune regulatory cytokine that acts on many cells of the immune system where it has profound anti-inflammatory functions, limiting excessive tissue disruption caused by inflammation. Mechanistically, IL10 binds to its heterotetrameric receptor comprising IL10RA and IL10RB leading to JAK1 and STAT2-mediated phosphorylation of STAT3. In turn, STAT3 translocates to the nucleus where it drives expression of anti-inflammatory mediators. Targets antigen-presenting cells (APCs) such as macrophages and monocytes and inhibits their release of pro-inflammatory cytokines including granulocyte-macrophage colony-stimulating factor /GM-CSF, granulocyte colony-stimulating factor /G-CSF, IL-1 alpha, IL-1 beta, IL-6, IL-8 and TNF-alpha. Interferes also with antigen presentation by reducing the expression of MHC-class II and co-stimulatory molecules, thereby inhibiting their ability to induce T cell activation.

For Research Use Only