# P53 Polyclonal Antibody

Catalog Number: E-AB-32467 2 Publications



Note: Centrifuge before opening to ensure complete recovery of vial contents.

	- 70	
Desc	$\mathbf{n}$	$\mathbf{n}$
DUSU.		

Reactivity Human, Monkey

**Immunogen** Synthesized peptide derived from human p53 around the non-phosphorylation site

of Ser392.

Host Rabbit Isotype IgG

**Purification** Affinity purification Conjugation **Unconjugated** 

**Formulation** PBS with 0.02% sodium azide, 0.5% protective protein and 50% glycerol, pH7.4

**Applications Recommended Dilution** 

WB 1:500-1:2000 **IHC** 1:100-1:300 IF 1:200-1:1000 **ELISA** 1:40000

#### Data



Western Blot analysis of COS7 cells using p53 Polyclonal Antibody at dilution of 1:2000.

> Observed Mw:47kDa Calculated Mw:44kDa

### **Preparation & Storage**

Storage Store at -20°C. Avoid freeze / thaw cycles.

## **Background**

Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclindependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. Implicated in Notch signaling cross-over. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis.

### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com