


Recombinant ROCK2+ROCK1 Monoclonal Antibody

catalog number: AN301910L

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description	
Reactivity	Human;Rat;Mouse
Immunogen	Recombinant human ROCK2+ROCK1 fragment
Host	Rabbit
Isotype	IgG, κ
Clone	C111531
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.
Applications	Recommended Dilution
WB	1:1000
IHC	1:50-1:100
IF	1:50


Data




Western Blot with ROCK2+ROCK1 Monoclonal Antibody at dilution of 1:1000. Lane 1: HeLa, Lane 2: Ramos, Lane 3: NIH/3T3, Lane 4: PC-12

Observed-MW:161 kDa


Calculated-MW:161, 158 kDa




Immunohistochemistry of paraffin-embedded Human colon cancer using ROCK2+ROCK1 Monoclonal Antibody at dilution of 1:100.



Immunohistochemistry of paraffin-embedded Mouse colon using ROCK2+ROCK1 Monoclonal Antibody at dilution of 1:100.



Immunohistochemistry of paraffin-embedded Rat cerebrum using ROCK2+ROCK1 Monoclonal Antibody at dilution of 1:100.



For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2

Immunofluorescent analysis of (4% Paraformaldehyde)
fixed HeLa cells using anti-ROCK2+ROCK1 Monoclonal
Antibody at dilution of 1:50.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

ROCK (Rho-associated kinase), a family of serine/threonine kinases, is an important downstream target of Rho-GTPase and plays an important role in Rho-mediated signaling. Two isoforms of ROCK have been identified: ROCK1 and ROCK2. ROCK is composed of N-terminal catalytic, coiled-coil, and C-terminal PH (pleckstrin homology) domains. The C-terminus of ROCK negatively regulates its kinase activity. ROCK1 is cleaved by caspase-3 at a conserved DETD1113/G sequence resulting in loss of its C-terminal inhibitory domain. ROCK2 is directly cleaved by granzyme B (grB). Cleavage activates ROCK2 and leads to phosphorylation of myosin light chain (MLC) and inhibition of myosin phosphatase. This phosphorylation may account for the mechanism by which Rho regulates cytokinesis, cell motility, cell membrane blebbing during apoptosis, and smooth muscle contraction.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2