

A Reliable Research Partner in Life Science and Medicine

## Recombinant ROCK2+ROCK1 Monoclonal Antibody

catalog number: AN301910L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human;Rat;Mouse

Immunogen Recombinant human ROCK2+ROCK1 fragment

Host Rabbit Isotype lgG, κ Clone A626


**Purification** Protein Apurified

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

#### **Applications Recommended Dilution**


1:1000 **WB** 1:50-1:100 **IHC** IF 1:50

#### Data




dilution of 1:1000. Lane 1: HeLa, Lane 2: Ramos, Lane 3:

NIH/3T3, Lane 4: PC-12 Observed-MW:161 kDa




Western Blot with ROCK2+ROCK1 Monoclonal Antibody at Immunohistochemistry of paraffin-embedded Human colon cancer using ROCK2+ROCK1 Monoclonal Antibody at dilution of 1:100.



Immunohistochemistry of paraffin-embedded Mouse colon using ROCK2+ROCK1 Monoclonal Antibody at dilution of

Immunohistochemistry of paraffin-embedded Rat cerebrum using ROCK2+ROCK1 Monoclonal Antibody at dilution of 1:100.

Rev. V1.1



### For Research Use Only

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

# **Elabscience**®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

Immunofluorescent analysis of (4% Paraformaldehyde) fixed HeLa cells using anti-ROCK2+ROCK1 Monoclonal Antibody at dilution of 1:50.

#### **Preparation & Storage**

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

#### **Background**

ROCK (Rho-associated kinase), a family of serine/threonine kinases, is an important downstream target of Rho-GTPase and plays an important role in Rho-mediated signaling. Two isoforms of ROCK have been identified: ROCK1 and ROCK2. ROCK is composed of N-terminal catalytic, coiled-coil, and C-terminal PH (pleckstrin homology) domains. The C-terminus of ROCK negatively regulates its kinase activity. ROCK1 is cleaved by caspase-3 at a conserved DETD1113/G sequence resulting in loss of its C-terminal inhibitory domain. ROCK2 is directly cleaved by granzyme B (grB). Cleavage activates ROCK2 and leads to phosphorylation of myosin light chain (MLC) and inhibition of myosin phosphatase. This phosphorylation may account for the mechanism by which Rho regulates cytokinesis, cell motility, cell membrane blebbing during apoptosis, and smooth muscle contraction.