NMT1 Polyclonal Antibody

Catalog Number: E-AB-68004



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human, Mouse, Rat

Immunogen Recombinant fusion protein of human NMT1

Host Rabbit
Isotype IgG


Purification Affinity purification
Conjugation Unconjugated

Formulation PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Applications Recommended Dilution

WB 1:200-1:2000

Data



Western blot analysis of extracts of various cell lines using NMT1 Polyclonal Antibody at 1:1000 dilution.

Observed MW:57kDa Calculated Mw:48kDa/56kDa

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Myristate, a rare 14-carbon saturated fatty acid, is cotranslationally attached by an amide linkage to the N-terminal glycine residue of cellular and viral proteins with diverse functions. N-myristoyltransferase (NMT; EC 2.3.1.97) catalyzes the transfer of myristate from CoA to proteins. N-myristoylation appears to be irreversible and is required for full expression of the biologic activities of several N-myristoylated proteins, including the alpha subunit of the signal-transducing guanine nucleotide-binding protein (G protein) GO (GNAO1; MIM 139311).

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com
Email: techsupport@elabscience.com