

A Reliable Research Partner in Life Science and Medicine

# PE Anti-Mouse CD40 Antibody [FGK4.5/FGK45]

Catalog Number: E-AB-F1028D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Mouse Host Rat

Isotype Rat IgG2a, κ
Clone No. FGK4.5/FGK45

**Isotype Control** PE Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832D]

Conjugation PE

**Conjugation Information** PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


## Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

## Data



Staining of C57BL/6 murine splenocytes with APC Anti-Mouse CD19 Antibody[1D3] and PE Anti-Mouse CD40 Antibody[FGK4.5/FGK45] (left) or PE Rat IgG2a, κ Isotype Control (right). Total viable cells were used for analysis.

## **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

## **Antigen Information**

Alternate Names B-cell surface antigen CD40;Bp50;CD40;CD40L receptor;Cd40;Tnfrsf5;Tumor necrosis

factor receptor superfamily member 5

Uniprot ID P27512

# For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

#### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

Gene ID Background 21939

CD40 is a 48 kD type I transmembrane glycoprotein also known as Bp50. It is a member of the tumor necrosis factor receptor (TNFR) superfamily and is expressed on B cells, basal epithelial cells, macrophages, follicular dendritic cells, endothelial cells, and a subset of CD34+ hematopoietic progenitors. CD40 regulates B cell developmen t/maturation, Ig isotype switching and, in combination with other signals such as IL-4, protects B cells from surface Ig-induced apoptosis and promotes proliferation. Interaction of CD40 with its ligand CD154 (gp39), which is expressed on activated T cells, is important in costimulation and immune regulation.