

Recombinant P4HB/ERBA2L Monoclonal Antibody

catalog number: **AN300349P**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human
Immunogen	Recombinant Human P4HB protein
Host	Rabbit
Isotype	IgG
Clone	10F8
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS

Applications Recommended Dilution

WB	1:500-1:2000
ICC/IF	1:20-1:100
IP	1-4 µL/mg of lysate

Data



Immunofluorescence analysis of Human P4HB in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 1% Triton X-100 in PBS, blocked with 10% serum, and incubated with Rabbit anti-Human P4HB monoclonal antibody (1:60) at 37°C 1 hour. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to endoplasmic reticulum.




Immunoprecipitation analysis using 2 µL anti-P4HB Monoclonal Antibody and 15 µl of 50 % Protein G agarose. Western blot was performed from the immunoprecipitate using P4HB Monoclonal Antibody at a dilution of 1:200.

Lane A: 0.5 mg MCF-7 Whole Cell Lysate

Observed-MW: 57 kDa

Calculated-MW: 57 kDa



Western Blot with P4HB Monoclonal Antibody at dilution of 1:100 dilution. Lane A: MCF7 Whole Cell Lysate, Lane B: HepG2 Whole Cell Lysate, Lysates/proteins at 30 µg per lane.

Observed-MW: 57 kDa

Calculated-MW: 57 kDa

Preparation & Storage

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.0

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping

Ice bag

Background

This gene encodes the beta subunit of prolyl 4-hydroxylase, a highly abundant multifunctional enzyme that belongs to the protein disulfide isomerase family. When present as a tetramer consisting of two alpha and two beta subunits, this enzyme is involved in hydroxylation of prolyl residues in procollagen. This enzyme is also a disulfide isomerase containing two thioredoxin domains that catalyze the formation, breakage and rearrangement of disulfide bonds. Other known functions include its ability to act as a chaperone that inhibits aggregation of misfolded proteins in a concentration-dependent manner, its ability to bind thyroid hormone, its role in both the influx and efflux of S-nitrosothiol-bound nitric oxide, and its function as a subunit of the microsomal triglyceride transfer protein complex.

For Research Use Only