

## Recombinant Mucin 5AC Monoclonal Antibody

catalog number: AN301037L

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


Reactivity	Human;Mouse;Rat
Immunogen	Recombinant Human Mucin 5AC protein
Host	Rabbit
Isotype	IgG,κ
Clone	B788
Purification	Protein A
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

### Applications


### Recommended Dilution

IHC	1:200-1:1000
WB	1:1000-1:5000
IF	1:200-1:1000
ELISA	1:5000-1:20000

### Data




Western Blot with Recombinant Mucin 5AC Monoclonal Antibody at dilution of 1:1000 dilution. Lane A: HT-29 cells.



Immunohistochemistry of paraffin-embedded human stomach tissue using Recombinant Mucin 5AC Monoclonal Antibody at dilution of 1:200.

**Observed-MW:586 kDa**

**Calculated-MW:586 kDa**



Immunohistochemistry of paraffin-embedded rat stomach tissue using Recombinant Mucin 5AC Monoclonal Antibody at dilution of 1:200.

### Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

### Background

#### For Research Use Only

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.2

The cysteine residues in the Cys-rich subdomain repeats are not involved in disulfide bonding.,Gel-forming glycoprotein of gastric and respiratory tract epithelia that protects the mucosa from infection and chemical damage by binding to inhaled microorganisms and particles that are subsequently removed by the mucociliary system.,PTM:C-, O- and N-glycosylated. O-glycosylated on the Thr-/Ser-rich tandem repeats. C-mannosylation in the Cys-rich subdomains may be required for proper folding of these regions and for export from the endoplasmic reticulum during biosynthesis.,PTM:Proteolytic cleavage in the C-terminal is initiated early in the secretory pathway and does not involve a serine protease. The extent of cleavage is increased in the acidic parts of the secretory pathway. Cleavage generates a reactive group which could link the protein to a primary amide.,similarity:Contains 1 CTCK (C-terminal cystine knot-like) domain.,similarity:Contains 2 WW domains.,similarity:Contains 4 WFD domains.,subunit: Multimeric. Interacts with H.pylori in the gastric epithelium, Barrett's esophagus as well as in gastric metaplasia of the duodenum (GMD).,tissue specificity:Highly expressed in surface mucosal cells of respiratory tract and stomach epithelia. Overexpressed in a number of carcinomas. Also expressed in Barrett's esophagus epithelium and in the proximal duodenum.

### For Research Use Only

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.2