

A Reliable Research Partner in Life Science and Medicine

Recombinant AID Monoclonal Antibody

catalog number: AN302047L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human;

Immunogen Peptide. This information is proprietary to PTMab.

 Host
 Rabbit

 Isotype
 IgG, κ


 Clone
 A767

Purification Protein Apurified

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution WB 1:1000 IHC 1:2000


Data



Western Blot with AID Monoclonal Antibody at dilution of 1:1000. Lane 1: K562(negative control), Lane 2: Ramos,

Immunohistochemistry of paraffin-embedded Human tonsil using AID Monoclonal Antibody at dilution of 1:2000.

Lane 3: Daudi
Observed-MW:23 kDa



Immunohistochemistry of paraffin-embedded Human diffuse large B-cell lymphoma using AID Monoclonal Antibody at dilution of 1:2000.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping lce bag

Background

For Research Use Only

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Activation-induced cytidine deaminase (AID) is thought to modify RNA due to its high homology to the RNA editing enzyme APOBEC-1. This function, however, has not been confirmed in in vitro studies, which show that AID has significant cytidine deaminase activity, and that this activity is blocked by zinc chelation. The B cell immune system must specifically recognize several infectious agents, which vastly outnumber immunoglobulin gene segments present in a given organism. Mechanisms such as somatic hypermutation, isotype switch recombination and gene conversion introduce diversity and specificity to the immune system. Analysis of mouse models and patients with AID deficiency has established a link between all three of these mechanisms and AID function. AID protein is detected in germinal center centroblast and germinal center derived lymphomas (Burkitt lymphoma), but not in pre-germinal center B cells or post-germinal center neoplasms (B cell chronic lymphocytic leukemia and multiple myeloma).

For Research Use Only