

Purified Anti-Mouse IL-2 Antibody[JES6-1A12], Functional Grade

catalog number: AN008330

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Immunogen	Recombinant Mouse IL-2 protein
Host	Rat
Isotype	Rat IgG2b, κ
Clone	JES6-1A12
Purification	>98%, Protein A/G purified
Buffer	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.

Applications

Recommended Dilution

FCM	$\leq 0.2 \mu\text{g}$ per million cells in 100 μL volume
Neut	Reported in the literature
Stim	Reported in the literature

Data



HEK293T cells transfected with pcDNA3.1 plasmid encoding Mouse IL-2 gene were stained with 0.2 μg Purified Anti-Mouse IL-2 Antibody[JES6-1A12], Functional Grade (Right) and 0.2 μg Rat IgG2b, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Rat IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.
Shipping	Ice bag

Background

Produced by T-cells in response to antigenic or mitogenic stimulation, this protein is required for T-cell proliferation and other activities crucial to regulation of the immune response. Can stimulate B-cells, monocytes, lymphokine-activated killer cells, natural killer cells, and glioma cells.

None (Azide-Free, Low Endotoxin) are perfectly suited to be used in culture or in vivo (for nonhuman studies) for functional assays blocking, neutralizing, activation or depletion where the presence of azide may damage cells or exogenous endotoxin may signal or activate cells.

For Research Use Only

Application References

K Kai McKinstry, et al. Nat Commun. 2014 Nov 5:5:5377. Alice McNally, et al. Proc Natl Acad Sci U S A. 2011 May 3; 108(18):7529-34.