

## Purified Anti-Mouse IL-2 Antibody[JES6-1A12], Functional Grade

catalog number: AN008330

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

### Description


<b>Reactivity</b>	Mouse
<b>Immunogen</b>	Recombinant MouseIL-2 protein
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG2b, κ
<b>Clone</b>	JES6-1A12
<b>Purification</b>	>98%, Protein A/G purified
<b>Buffer</b>	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.

### Applications

### Recommended Dilution

<b>FCM</b>	2 μg/mL(0.5×10 <sup>6</sup> -1×10 <sup>6</sup> cells)
<b>Neut</b>	Reported in the literature
<b>Stim</b>	Reported in the literature

### Data



HEK293T cells transfected with pcDNA3.1 plasmid encoding Mouse IL-2 gene were stained with 0.2 μg Purified Anti-Mouse IL-2 Antibody[JES6-1A12], Functional Grade (Right) and 0.2 μg Rat IgG2b, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Rat IgG Secondary Antibody.

### Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.
<b>Shipping</b>	Ice bag

### Background

Produced by T-cells in response to antigenic or mitogenic stimulation, this protein is required for T-cell proliferation and other activities crucial to regulation of the immune response. Can stimulate B-cells, monocytes, lymphokine-activated killer cells, natural killer cells, and glioma cells.

None (Azide-Free, Low Endotoxin) are perfectly suited to be used in culture or in vivo (for nonhuman studies) for functional assays blocking, neutralizing, activation or depletion where the presence of azide may damage cells or exogenous endotoxin may signal or activate cells.

### For Research Use Only

## Application References

K Kai McKinstry, et al. Nat Commun. 2014 Nov 5:5:5377. Alice McNally, et al. Proc Natl Acad Sci U S A. 2011 May 3; 108(18):7529-34.