

IFITM3 Polyclonal Antibody

catalog number: D-AB-10162L

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Mouse;Rat
Immunogen	Recombinant Rat IFITM3 protein expressed by E.coli
Host	Rabbit
Isotype	IgG
Purification	Antigen Affinity Purification
Buffer	PBS with 0.05% Proclin300, 1% protective protein and 50% glycerol, pH7.4

Applications

Applications	Recommended Dilution
WB	1:1000-1:2000
IHC	1:50-1:200
IF	1:100-1:400

Data




Western blot with IFITM3 Polyclonal antibody at dilution of 1:2000. lane 1: Hela whole cell lysate, lane 2: 293 T whole cell lysate, lane 3: Mouse Liver, lane 4: Mouse Heart, lane 5: Rat



Liver

Observed-MW: 15 kDa


Calculated-MW: 15 kDa




Immunohistochemistry of paraffin-embedded Mouse lung using IFITM3 Polyclonal Antibody at dilution of 1:100



Immunohistochemistry of paraffin-embedded Rat kidney using IFITM3 Polyclonal Antibody at dilution of 1:100



Immunohistochemistry of paraffin-embedded Human lung cancer using IFITM3 Polyclonal Antibody at dilution of 1:200



For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V2.1

Immunofluorescence analysis of Hela cells using IFITM3

Immunofluorescence analysis of Hela cells using ifitm3

Polyclonal Antibody at dilution of 1:100

Polyclonal Antibody at dilution of 1:100

Preparation & Storage

Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping

The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

IFITM3, also named as interferon-inducible protein 1-8U, belongs to the CD225 family. It is IFN-induced antiviral protein that mediates cellular innate immunity to at least three major human pathogens, namely influenza A H1N1 virus, West Nile virus (WNV), and dengue virus, by inhibiting the early steps of replication. IFITM3 is identified as interferon-induced cellular proteins that restrict infections by retroviruses and filoviruses and of influenza virus and flaviviruses, respectively. IFITM3, the most potent antiviral IFITM, was found to inhibit an uncharacterized early infectious event after VSV endocytosis, but before primary transcription of its viral genome. IFITM proteins are viral restriction factors that can inhibit infection mediated by the influenza A virus (IAV) hemagglutinin (HA) protein. They differentially restrict the entry of a broad range of enveloped viruses, and modulate cellular tropism independently of viral receptor expression. This antibody recognizes both IFITM2 and IFITM3.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V2.1