# Recombinant Rat CD147/BSG/Basigin Protein (His Tag)

Catalog Number: PDER100147



Note: Centrifuge before opening to ensure complete recovery of vial contents.

|     |              | crip |    |   |    |    |
|-----|--------------|------|----|---|----|----|
|     | 00           | CI   | РΤ | n | tπ | Λn |
| JU. | $\mathbf{c}$ | v.   |    | w | ш  | UП |

Species Rat

Source E.coli-derived Rat CD147 protein Pro52-Thr324, with an N-terminal His

 Mol\_Mass
 30.3 kDa

 Accession
 P26453-1

**Bio-activity** Not validated for activity

#### **Properties**

**Purity** > 95% as determined by reducing SDS-PAGE.

Endotoxin < 10 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

**Reconstitution** It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



> 95 % as determined by reducing SDS-PAGE.

#### Background

BSG,also named as 5F7,Basigin,EMMPRIN,TCSF and CD147,plays pivotal roles in spermatogenesis,embryo implantatio n,neural network formation and tumor progression. It stimulates adjacent fibroblasts to produce matrix metalloproteinases (MMPS).CD147 may target monocarboxylate transporters SLC16A1,SLC16A3 and SLC16A8 to plasma membranes of retinal pigment epithelium and neural retina. It seems to be a receptor for oligomannosidic glycans. CD147 is a receptor of CypA,inducing matrix metalloproteinase expression and mediating the degradation of the extracellular matrix,plays an important role in tumorigenesis and invasion in oral cancer. It has been considered as an objective and effective marker to predict invasion and prognosis in various cancers.

## For Research Use Only