

## Recombinant Human RELA/Transcription factor p65/NFkB p65 Protein (aa 1-306, GST Tag)

Catalog Number: PKSH030921

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human RELA/Transcription factor p65/NFkB p65 protein Met 1-Tyr 306, with an N-terminal GST
<b>Calculated MW</b>	62.0 kDa
<b>Observed MW</b>	58 kDa
<b>Accession</b>	Q04206-1
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 85 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 20mM Tris, 0.15M NaCl, 20mM GST, pH 8.0 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 85 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

RELA (v-rel reticuloendotheliosis viral oncogene homolog A), also known as Nuclear factor NF-kappa-B p65 subunit, or Transcription factor p65, is a transcription factor expressed in growth plate chondrocytes where it facilitates chondrogenesis. The v-rel avian reticuloendotheliosis viral oncogene homolog A (RELA) gene encodes the major component of the NF-κB complex. NF-kappaB is a generic name for an evolutionarily conserved transcription-factor system that contributes to the mounting of an effective immune response but is also involved in the regulation of cell proliferation, development, and apoptosis. The implication of NF-kappaB in central biological processes and its extraordinary connectivity to other signaling pathways raise a need for highly controlled regulation of NF-kappaB activity at several levels. The mammalian Rel/NF-kappaB family of transcription factors, including RelA, c-Rel, RelB, NF-kappaB1 (p50 and its precursor p105), and NF-kappaB2 (p52 and its precursor p100), plays a central role in the immune system by regulating several processes ranging from the development and survival of lymphocytes and lymphoid organs to the control of immune responses and malignant transformation.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017