

## Purified Anti-Human CD123 Antibody[HI8D7]

catalog number: **AN008000P**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human
<b>Immunogen</b>	Recombinant Human CD123 protein
<b>Host</b>	Mouse
<b>Isotype</b>	Mouse IgG1, $\kappa$
<b>Clone</b>	HI8D7
<b>Purification</b>	>98%, Protein A/G purified
<b>Buffer</b>	PBS, pH 7.2. Contains 0.05% proclin 300.

### Applications

### Recommended Dilution

<b>FCM</b>	2 $\mu$ g/mL( $1 \times 10^5$ - $5 \times 10^5$ cells)
------------	--

### Data



Human peripheral blood lymphocytes cell were stained with 0.2  $\mu$ g Purified Anti-Human CD123 Antibody[HI8D7] (Right) and 0.2  $\mu$ g Mouse IgG1,  $\kappa$  Isotype Control (Left), followed by

Elab Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD45RO FITC-conjugated Monoclonal Antibody.

### Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	Ice bag

### Background

CD123 is the 70 kD transmembrane  $\alpha$  chain of the IL-3 receptor. Alone, CD123 binds IL-3 with low affinity; when CD123 associates with CDw131 (common  $\beta$  chain), it binds IL-3 with high affinity. CD123 does not transduce intracellular signals upon binding IL-3 and requires the  $\beta$  chain for this function. CD123 is expressed by myeloid precursors, macrophages, dendritic cells, mast cells, basophils, megakaryocytes, and some B cells.

### For Research Use Only