

## Recombinant Cynomolgus OX40 Ligand/TNFSF4/OX40L Protein (His Tag)

Catalog Number: PKSQ050028

Note: Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Cynomolgus macaques
<b>Source</b>	HEK293 Cells-derived Cynomolgus macaques TNFSF4/OX40L protein Gln51-Leu183, with an N-terminal His
<b>Calculated MW</b>	16.5 kDa
<b>Observed MW</b>	23-30 kDa
<b>Accession</b>	F7FL80
<b>Bio-activity</b>	Immobilized Human OX40-Fc(PKSH032843) at 10µg/ml (100 µl/well) can bind Cynomolgus OX40L-His(PKSQ050028). The ED <sub>50</sub> of Cynomolgus OX40L-His(PKSQ050028) is 17.75 ng/ml.


### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.



Immobilized Human OX40-Fc(PKSH032843) at 10µg/ml (100 µl/well) can bind Cynomolgus OX40L-His(PKSQ050028). The ED<sub>50</sub> of Cynomolgus OX40L-His(PKSQ050028) is 17.75 ng/ml.

### Background

#### For Research Use Only

Tumor necrosis factor ligand superfamily member 4 (TNFSF4/OX40L) is a single-pass type II membrane protein. OX40L is expressed by DCs, macrophages and B cells and signals via its cognate receptor OX40 which is mainly expressed on APCs. OX40L/OX40 interactions are important in T-cell activation and survival and for the generation of memory T cells from activated effector T cells. It is a cytokine that binds to TNFRSF4, Co-stimulates T-cell proliferation and cytokine production. OX40L/OX40 interactions are important in T-cell activation and survival and for the generation of memory T cells from activated effector T cells.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017