



EIF4A2 Polyclonal Antibody

catalog number: E-AB-19148

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Human;Mouse;Rat |
| Immunogen | Fusion protein of human EIF4A2 |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Antigen affinity purification |
| Conjugation | Unconjugated |
| Buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution |
| IHC | 1:50-1:200 |

| Data | |
|--|--|
|  |  |
| Immunohistochemistry of paraffin-embedded Human liver cancer tissue using EIF4A2 Polyclonal Antibody at dilution of 1:65(×200) | Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using EIF4A2 Polyclonal Antibody at dilution of 1:65(×200) |

| Preparation & Storage | |
|-----------------------|--|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended. |

| Background | |
|---|--|
| ATP-dependent RNA helicase which is a subunit of the eIF4F complex involved in cap recognition and is required for mRNA binding to ribosome. In the current model of translation initiation, eIF4A unwinds RNA secondary structures in the 5'-UTR of mRNAs which is necessary to allow efficient binding of the small ribosomal subunit, and subsequent scanning for the initiator codon. | |