

Recombinant HMGB2 Monoclonal Antibody

catalog number: AN301821L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Rat; Mouse

Immunogen Recombinant human HMGB2 fragment


HostRabbitIsotypeIgG, κCloneA533


Purification Protein A purified

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications	Recommended Dilution
WB	1:500-1:1000
IHC	1:50-1:100
IF	1:50
IP	1:25-1:50


Data







Western Blot with HMGB2 Monoclonal Antibody at dilution of Immunohistochemistry of paraffin-embedded Human colon 1:1000. Lane 1: HeLa, Lane 2: HepG2, Lane 3: PC-12 using HMGB2 Monoclonal Antibody at dilution of 1:100.

Observed-MW:24 kDa Calculated-MW:24 kDa






Immunohistochemistry of paraffin-embedded Mouse testis using HMGB2 Monoclonal Antibody at dilution of 1:100.



Immunohistochemistry of paraffin-embedded Rat cerebrum using HMGB2 Monoclonal Antibody at dilution of 1:100.



For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Immunofluorescent analysis of (4% Paraformaldehyde) fixed Immunoprecipitation analysis using anti-HMGB2 Monoclonal HeLa cells using anti-HMGB2 Monoclonal Antibody at Antibody. Western blot was performed from the immunoprecipitate using HMGB2 Monoclonal Antibody at a dilution of 1:50. Lane 1: 10% Input, Lane 2: HMGB2

Isotype
Observed-MW:24 kDa
Calculated-MW:24 kDa

Monoclonal Antibody, Lane 3: Rabbit monoclonal IgG

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

HMGB2 belongs to a family of highly conserved proteins that contain HMG box domains. HMGB2 is widely expressed during embryonic development, but it is restricted to lymphoid organs and testis in adult animals. HMGB2 facilitate the binding of Hox proteins, Oct proteins, p53, Rel proteins, and steroid hormone receptor proteins to their target gene promoters. Furthermore, HMGB2 interacts with RAG1 to facilitate RAG complex binding to the recombinant signal sequence (RSS) and stimulate DNA-bending and subsequent VDJ cleavage at antigen receptor genes. In addition to their functions in the nucleus, HMGB proteins play a significant role in extracellular signaling associated with inflammation. HMGB2 is secreted by myeloid cells and promotes proliferation and migration of endothelial cells by binding to the receptor for advanced glycation end products (RAGE).

For Research Use Only