Elabscience®

SLC1A6 Polyclonal Antibody

catalog number: E-AB-15608

Note: Centrifuge before opening to ensure complete recovery of vial contents.

1:50-1:200

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Synthetic peptide of human SLC1A6
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000

Data

IHC





cancer using SLC1A6 Polyclonal Antibody at dilution of

1:40

Western Blot analysis of Human testis tissue and A375 cell using SLC1A6 Polyclonal Antibody at dilution of 1:400

Calculated-MW:62 kDa



Immunohistochemistry of paraffin-embedded Human colon

cancer using SLC1A6 Polyclonal Antibody at dilution of 1.40

cancer using SECTAO Toryclonar Antibody at unution of 1.40	
Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017

Elabscience®

Excitatory Amino Acid Transporters (EAATs) are membrane-bound proteins that are localized in glial cells and presynaptic glutamatergic nerve endings. EAATs transport the excitatory neurotransmitters L-glutamate and D-aspartate, a process that is essential for terminating the postsynaptic action of glutamate. The re-uptake of amino acid neurotransmitters by EAAT proteins has been shown to protect neurons from excitotoxicity, which is caused by the accumulation of amino acid neurotransmitters. EAAT4 is an aspartate/glutamate transporter that is expressed predominantly in the cerebellum. The transport activity encoded by EAAT4 has high apparent affinity for L-aspartate and L-glutamate, and has a pharmacologic profile consistent with previously described cerebellar transport activities. EAAT5 is a glutamate transporter coupled to a chloride conductance which is expressed primarily in retina. Although EAAT5 shares the structural homologies of the EAAT family, a novel feature of the EAAT5 sequence is a carboxyterminal motif previously identified in N-ethyl-D-aspartate receptors and potassium channels and shown to confer interactions with a family of synaptic proteins that promote ion channel clustering.

For Research Use Only