

SHTN1 Polyclonal Antibody

catalog number: **E-AB-18189**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description



Reactivity	Human
Immunogen	Synthetic peptide of human SHTN1
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Recommended Dilution

WB	1:500-1:2000
IHC	1:50-1:100

Data



Western blot analysis of Human fetal liver tissue lysate using SHTN1 Polyclonal Antibody at dilution of 1:600

Immunohistochemistry of paraffin-embedded Human tonsil tissue using SHTN1 Polyclonal Antibody at dilution of 1:60(x200)

Observed-MW:Refer to figures

1:60(x200)

Calculated-MW:72 kDa

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

Involved in the generation of internal asymmetric signals required for neuronal polarization and neurite outgrowth. Mediates netrin-1-induced F-actin-substrate coupling or 'clutch engagement' within the axon growth cone through activation of CDC42, RAC1 and PAK1-dependent signaling pathway, thereby converting the F-actin retrograde flow into traction forces, concomitantly with filopodium extension and axon outgrowth. Plays a role in cytoskeletal organization by regulating the subcellular localization of phosphoinositide 3-kinase (PI3K) activity at the axonal growth cone. Plays also a role in regenerative neurite outgrowth. In the developing cortex, cooperates with KIF20B to promote both the transition from the multipolar to the bipolar stage and the radial migration of cortical neurons from the ventricular zone toward the superficial layer of the neocortex. Involved in the accumulation of phosphatidylinositol 3,4,5-trisphosphate (PIP3) in the growth cone of primary hippocampal neurons.

For Research Use Only