

Recombinant Human DPEP2 Protein (Fc Tag)

Catalog Number: PKSH030555

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Human
Source	HEK293 Cells-derived Human DPEP2 protein Met 1-Ser376, with an C-terminal hFc
Calculated MW	65.0 kDa
Observed MW	43-49 kDa
Accession	AAH24021.1
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

DPEP2 (MBD-2) belongs to the membrane-bound dipeptidase family. There are three members of this family as membrane-bound dipeptidase-1 (MBD-1), membrane-bound dipeptidase-2 (MBD-2) and membrane-bound dipeptidase-3 (MBD-3). MBD-2 is expressed at highest levels in lung, heart, and testis and at some what lower levels in spleen. MBD-2 is membrane-bound through a glycosylphosphatidyl-inositol linkage and probably is a metalloprotease which hydrolyzes leukotriene D4 (LTD4) into leukotriene E4 (LTE4). It is generally recognized that rapid cleavage of LTD4 is important in inactivating the broncho- and vaso-constrictive effects of cysteinyl LTs in asthmatic and inflammatory processes.

For Research Use Only