Recombinant Human DPEP2 Protein (Fc Tag)

Catalog Number: PKSH030555

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Human
HEK293 Cells-derived Human DPEP2 protein Met 1-Ser376, with an C-terminal hFc
65.0 kDa
43-49 kDa
AAH24021.1
Not validated for activity
>95 % as determined by reducing SDS-PAGE.
< 1.0 EU per µg of the protein as determined by the LAL method.
Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
This product is provided as lyophilized powder which is shipped with ice packs.
Lyophilized from sterile PBS, pH 7.4
Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
before lyophilization.
Please refer to the specific buffer information in the printed manual.
Please refer to the printed manual for detailed information.

Data

KDa 116	MK	R
66.2	-	
45.0		
35.0	-	
25.0	-	
18.4 14.4	=	

> 95 % as determined by reducing SDS-PAGE.

Background

DPEP2 (MBD-2) belongs to the membrane-bound dipeptidase family. There are three members of this family as membran e-bound dipeptidase-1 (MBD-1),membrane-bound dipeptidase-2 (MBD-2) and membrane-bound dipeptidase-3 (MBD-3). MBD-2 is expressed at highest levels in lung, heart, and testis and at some what lower levels in spleen.MBD-2 is membrane-bound through a glycosylphosphatidyl-inositol linkage and probably is a metalloprotease which hydrolyzes leukotriene D4 (LTD4) into leukotriene E4 (LTE4). It is generally recognized that rapid cleavage of LTD4 is important in inactivating the broncho-and vaso-constrictive effects of cysteinyl LTs in asthmatic and inflammatory processes.

Web:www.elabscience.com