

# Recombinant Mouse Ccl7 Protein(GST Tag)

Catalog Number: PDEM100323

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Species</b>	Mouse
<b>Source</b>	E.coli-derived Mouse Ccl7 protein Gln24-Pro97, with an N-terminal GST
<b>Mol_Mass</b>	34.1 kDa
<b>Accession</b>	Q03366
<b>Bio-activity</b>	Not validated for activity

## Properties

<b>Purity</b>	> 95% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 10 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis

## Data



SDS-PAGE analysis of Mouse Ccl7 proteins, 2µg/lane of Recombinant Mouse Ccl7 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 39 kDa

## Background

### For Research Use Only

# Recombinant Mouse Ccl7 Protein(GST Tag)

Catalog Number: PDEM100323



Chemokines are a family of small chemotactic cytokines, or proteins secreted by cells. Chemokines share the same structure similarities such as small size, and the presence of four cysteine residues in conserved locations in order to form their 3-dimensional shape. Some of the chemokines are considered pro-inflammatory which can be induced to recruit cells of the immune system to a site of infection during an immune response, while others are considered homeostatic and are implied in controlling the migration of cells during normal processes of tissue maintenance and development. There are four members of the chemokine family: C-C kemokines, C kemokines, CXC kemokines and CX3C kemokines. The C-C kemokines have two cysteines nearby the amino terminus. There have been at least 27 distinct members of this subgroup reported for mammals, called C-C chemokine ligands-1 to 28. Chemokine ligand 7(CCL7), also known as MCP-3, is a isform of the C-C chemokine subfamily of the chemokine family which is produced by certain tumor cells and by macrophages. It also own two adjacent N-terminal cysteine residues. Chemokine ligand 7(CCL7) spacifically attracts monocytes, and regulates macrophage function.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine  
Tel:400-999-2100

Email:[techsupport@elabscience.cn](mailto:techsupport@elabscience.cn)

Web:[www.elabscience.cn](http://www.elabscience.cn)

Rev. V1.0