

Recombinant Rat FGF-2/FGFb Protein

Catalog Number: PKSR030423

Note: Centrifuge before opening to ensure complete recovery of vial contents.



Description

Species	Rat
Source	E.coli-derived Rat FGF-2/FGFb protein Ala11-Ser154
Calculated MW	16.2 kDa
Observed MW	16 kDa
Accession	P13109
Bio-activity	Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells. The ED ₅₀ for this effect is 0.3-1. 8 ng/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells. The ED₅₀ for this effect is 0.3-1.8 ng/ml.

Background

For Research Use Only

FGF-basic is a members of the Fibroblast Growth Factors (FGFs) family. The family constitutes a large family of proteins involved in many aspects of development including cell proliferation, growth, and differentiation. They act on several cell types to regulate diverse physiologic functions including angiogenesis, cell growth, pattern formation, embryonic development, metabolic regulation, cell migration, neurotrophic effects, and tissue repair. FGF-basic is a non-glycosylated heparin binding growth factor that is expressed in the brain, pituitary, kidney, retina, bone, testis, adrenal gland liver, monocytes, epithelial cells and endothelial cells. FGF-basic signals through FGFR 1b, 1c, 2c, 3c and 4.