



A Reliable Research Partner in Life Science and Medicine

Recombinant Human MAP4K5 Protein (His Tag)

Catalog Number: PDEH101049

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source E.coli-derived Human MAP4K5 protein Phe601-Tyr846, with an N-terminal His & C-

terminal His

 Calculated MW
 27.0 kDa

 Observed MW
 35 kDa

 Accession
 Q9Y4K4

Bio-activity Not validated for activity

Properties

Purity > 95% as determined by reducing SDS-PAGE.

Endotoxin < 10 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping

This product is provided as lyophilized powder which is shipped with ice packs.

Formulation

Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

Background

Mitogen-activated protein kinase kinase kinase kinase 5, also known as Kinase homologous to SPS1/STE20, MAPK/ ERK kinase kinase 5, MEK kinase kinase 5 and MAP4K5, is a cytoplasm protein which belongs to the , protein kinase superfamily, STE Ser/Thr protein kinase family and STE20 subfamily. MAP4K5 is ubiquitously expressed in all tissues examined, with high levels in the ovary, testis and prostate. It contains one ,CNH domain and one ,protein kinase domain. MAP4K5 is highly similar to yeast SPS1/STE20 kinase. Yeast SPS1/STE20 functions near the beginning of the MAP kinase signal cascades that is essential for yeast pheromone response. MAP4K5 has been shown to interact with CRKL and TRAF2. This kinase was shown to activate Jun kinase in mammalian cells. MAP4K5 is an early component of MAP kinase signal cascades. It may play a role in the response to environmental stress. MAP4K5 appears to act upstream of the JUN N-terminal pathway.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com