FADD Polyclonal Antibody

catalog number: **E-AB-93254**


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of mouse FADD
Host	Rabbit
Isotype	Rabbit IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.


Applications	Recommended Dilution
WB	1:200-1:2000
IF	1:50-1:200
IP	1:50-1:200

Data




Western blot analysis of extracts of Mouse heart using FADD Polyclonal Antibody at 1:1000 dilution.

Observed-MW:28 kDa




Western blot analysis of extracts of Jurkat cells using FADD Polyclonal Antibody at 1:1000 dilution.

Observed-MW:28 kDa



Immunofluorescence analysis of mouse bone marrow cells using FADD Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunoprecipitation analysis of 600ug extracts of Mouse heart cells using 3ug FADD Polyclonal Antibody. Western blot was performed from the immunoprecipitate using FADD Polyclonal Antibody at a dilution of 1:1000.

Observed-MW:28 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com

Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95 or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling.