
Elabscience®

Elab Fluor[®] Violet 610 Anti-Mouse CD11c Antibody[N418]

Catalog Number: E-AB-F0991T

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Armenian Hamster
lsotype	Armenian Hamster IgG
Clone No.	N418
Isotype Control	Elab Fluor [®] Violet 610 Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09852T]
Conjugation	Elab Fluor [®] Violet 610
Conjugation Information	Elab Fluor [®] Violet 610 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 613 nm (e.g., a 615/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.



Staining of C57BL/6 murine splenocytes with FITC Anti-

Mouse MHC II (I-A/I-E) Antibody[M5/114] and Elab Fluor[®] Violet 610 Anti-Mouse CD11c Antibody[N418] (left) or Elab

Fluor[®] Violet 610 Armenian Hamster IgG Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storag	e
Storage	Keep as concentrated solution.
.	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	lce bag
Antigen Information	
Alternate Names	CD11 antigen-like family member C;CD11c;Integrin alpha-X;Itgax;Leukocyte adhesion receptor p150+95
For Research Use C	Dnly

Elabscience®

Uniprot ID Gene ID Background Q9QXH4

16411

CD11c is a 150 kD glycoprotein also known as α X integrin, CR4, and p150. CD11c forms a α X β 2 heterodimer with β 2 integrin (CD18). It is primarily expressed on dendritic cells, NK cells, a subset of intestinal intraepithelial lymphocytes (IEL), and some activated T cells. The α X β 2 integrin plays an important role in cell-cell contact by binding its ligands: iC3b, fibrinogen and CD54.