

## Mouse HGF Antibody Pair Set

<b>Catalog No.</b>	E-KAB-0338	<b>Applications</b>	ELISA
<b>Synonyms</b>	F-TCF, HGFB, HPTA, SF, Scatter Factor, Hepapoietin A		

### Kit components & Storage

Title	Specifications	Storage
Mouse HGF Capture Antibody	1 vial, 100 µg	Store at -20℃ for one year. Avoid freeze / thaw cycles.
Mouse HGF Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20℃ for one year. Avoid freeze / thaw cycles.

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Product Information

Items		Characteristic (E-KAB-0338)	
		Mouse HGF Capture Antibody	Mouse HGF Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Mouse HGF protein	Recombinant Mouse HGF protein
	Swissprot	Q08048	
Product details	Reactivity	Mouse	Mouse
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Mouse HGF in ELISAs.	

### For Research Use Only

## Applications

### Mouse HGF Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images																		
ELISA Capture	0.5-4μg/mL	Mouse HGF Capture Antibody	 <table><caption>Approximate data points from the standard curve</caption><thead><tr><th>Mouse HGF concentration (pg/mL)</th><th>Optical Density</th></tr></thead><tbody><tr><td>10</td><td>0.05</td></tr><tr><td>20</td><td>0.1</td></tr><tr><td>50</td><td>0.2</td></tr><tr><td>100</td><td>0.4</td></tr><tr><td>200</td><td>0.8</td></tr><tr><td>500</td><td>1.5</td></tr><tr><td>1000</td><td>2.5</td></tr><tr><td>2000</td><td>4.0</td></tr></tbody></table>	Mouse HGF concentration (pg/mL)	Optical Density	10	0.05	20	0.1	50	0.2	100	0.4	200	0.8	500	1.5	1000	2.5	2000	4.0
Mouse HGF concentration (pg/mL)	Optical Density																				
10	0.05																				
20	0.1																				
50	0.2																				
100	0.4																				
200	0.8																				
500	1.5																				
1000	2.5																				
2000	4.0																				
ELISA Detection	1:1000-1:10000	Mouse HGF Detection Antibody (Biotin)																			

**Note:** This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

## Background

Hepatocyte growth factor regulates cell growth, cell motility, and morphogenesis by activating a tyrosine kinase signaling cascade after binding to the proto-oncogenic c-Met receptor. Hepatocyte growth factor is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. Its ability to stimulate mitogenesis, cell motility, and matrix invasion gives it a central role in angiogenesis, tumorigenesis, and tissue regeneration. It is secreted as a single inactive polypeptide and is cleaved by serine proteases into a 69-kDa alpha-chain and 34-kDa beta-chain. A disulfide bond between the alpha and beta chains produces the active, heterodimeric molecule.