

## Recombinant Human DCBLD2/ESDN Protein (aa 1-482, His Tag)

**Catalog Number:** PKSH030592

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human DCBLD2/ESDN protein Met 1-Ala482, with an C-terminal His
<b>Calculated MW</b>	52.6 kDa
<b>Observed MW</b>	69-89 kDa
<b>Accession</b>	NP_563615.3
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

DCBLD2, also known as ESDN and CLCP1, localizes in various compartments. DCBLD2 is up-regulated in lung cancers and is regulated by transcription factor AP-2 alpha (TFAP2A), a component of activator protein-2 (AP-2) that is known to regulate IL-8 production in human lung fibroblasts and epithelial cells. DCBLD2 could be related to FEV(1)-related phenotypes in asthmatics. DCBLD2 gene is expressed at very high level. DCBLD2 is proposed to participate in processes such as intracellular receptor mediated signaling pathway, negative regulation of cell growth and so on.

### For Research Use Only