## Phospho-lkB alpha (Ser32/S36) Polyclonal Antibody

## catalog number: E-AB-20911

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description  |   |
|--|---|
| Reactivity   | Human;Mouse;Rat;Monkey  |
| Immunogen  | Synthesized peptide derived from human I $\kappa$ B- $\alpha$ around the phosphorylation site of Ser32/36   |
| Host   | Rabbit  |
| Isotype  | IgG   |
| Purification   | Affinity purification   |
| Buffer   | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein protectant and 50% glycerol. |
| Applications   | Recommended Dilution  |
| WB   | 1:500-1:2000  |
| IHC  | 1:100-1:300   |
| IF   | 1:200-1:1000  |
| Data   |   |
| alpha (Ser32)<br>Obse<br>Calcu<br>Immunofluorescence<br>Phospho-IkB alpha (S | <ul> <li>analysis of Rat spleen tissue using ber32/S36) Polyclonal Antibody at bition of 1:200</li> </ul>   |
|  | lution of 1:200   |
| Preparation & Storage  |   |
| Storage  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.   |
| Shipping   | The product is shipped with ice pack, upon receipt, store it immediately at the                             |

Background

## For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

## **Elabscience**®

Inhibits the activity of dimeric NF-kappa-B/REL complexes by trapping REL dimers in the cytoplasm through masking of their nuclear localization signals. On cellular stimulation by immune and proinflammatory responses, becomes phosphorylated promoting ubiquitination and degradation, enabling the dimeric RELA to translocate to the nucleus and activate transcription.

For Research Use Only