

Recombinant Human Glypican 5/GPC5 Protein (His Tag)

Catalog Number: PKSH031890

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	Baculovirus-Insect Cells-derived Human Glypican 5/GPC5 protein Met 1-Thr 554, with an C-terminal His
Calculated MW	60.5 kDa
Observed MW	60.5 kDa
Accession	NP_004457.1
Bio-activity	Immobilized human GPC5 at 5 µg/ml (100 µl/well) can bind human bFGF with a linear ranger of 0.156-2.5 ng/ml.

Properties

Purity	> 92 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 50mM Tris, 100mM NaCl, pH 8.0 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



Measured by its binding ability in a functional ELISA.
Immobilized human GPC5 at 5 µg/ml (100 µl/well) can bind human bFGF with a linear range of 0.156-2.5 ng/ml.

Background

For Research Use Only

Glypican-5 (GPC5), is a cell membrane protein which belongs to the glypican family. The glypicans compose a family of glycosylphosphatidylinositol-anchored heparan sulfate proteoglycans that may play a role in the control of cell division and growth regulation. So far, six members (Glypican-1/GPC1, Glypican-2/GPC2, Glypican-3/GPC3, Glypican-4/GPC4, Glypican-5/GPC5, Glypican-6/GPC6) of this family are known in vertebrates. In adult, Glypican-5 is primarily expressed in the brain. It is also detected in fetal brain, lung and liver. Glypican-5 enhances the intracellular signaling of FGF2 and HGF. It alters the cellular distribution of FGF2. The properties of Glypican-5 make it an attractive target for therapeutic intervention in rhabdomyosarcomas and other tumors that amplify and/or overexpress its gene. Glypican-5 is over-expressed in lymphoma cell lines that had shown amplification. It is a likely target for amplification, and that over-expression of GPC5 may contribute to development and/or progression of lymphomas and other tumors.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017