



A Reliable Research Partner in Life Science and Medicine

Recombinant HIP2/LIG Monoclonal Antibody

catalog number: AN301704L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human;

Immunogen Recombinant human HIP2/LIG fragment

HostRabbitIsotype IgG, κ CloneA412


Purification Protein A purified

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution

WB 1:500-1:2000

Data



Western Blot with HIP2/LIG Monoclonal Antibody at dilution

of 1:2000. Lane 1: HeLa, Lane 2: 293T, Lane 3: HepG2,

Lane 4: PC-3
Observed-MW:22 kDa
Calculated-MW:22 kDa

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

Huntingtin-interacting protein 2 (HIP2/LIG), is a E2 ubiquitin-conjugating enzyme, which accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In vitro, HIP2/LIG catalyzes the synthesis of 'Lys-48'-linked polyubiquitin chains in the presence or in the absence of BRCA1-BARD1 E3 ubiquitin-protein ligase complex, while it does not transfer ubiquitin directly to but elongates monoubiquitinated substrate protein. Moreover, HIP2/LIG mediates the selective degradation of short-lived and abnormal proteins, such as the endoplasmic reticulum-associated degradation (ERAD) of misfolded lumenal proteins.

For Research Use Only