## Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

# Rat IgG2c Antibody Pair Set

Catalog No. E-KAB-0632 Applications ELISA

Synonyms IgG2c

## Kit components & Storage

Title	Specifications	Storage
Rat IgG2c Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze/thaw cycles.
Rat IgG2c Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20°C for one year.
		Avoid freeze/thaw cycles.

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0632)		
		Rat IgG2c Capture Antibody	Rat IgG2c Detection Antibody (Biotin)	
Immunogen	Immunogen	Natural Rat IgG2c protien	Natural Rat IgG2c protien	
Information	Swissprot	/		
Product details	Reactivity	Rat	Rat	
	Host	Mouse	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5 mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%	
		glycerol; pH 7.5	protective protein; 50% glycerol; pH	
			7.5	
	Purify	Protein A or G	Antigen Affinity	
Specificity		Detects Rat IgG2c in ELISAs.		

For Research Use Only

Tel: 400-999-2100 Web: <a href="www.elabscience.cn">www.elabscience.cn</a> Email: <a href="techsupport@elabscience.cn">techsupport@elabscience.cn</a>

## Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

## **Applications**

Rat IgG2c Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Rat IgG2c Capture	
Capture		Antibody	10
			Ajsu
ELISA	1:1000-1:10000	Rat IgG2c Detection	Optical Density
Detection		Antibody (Biotin)	Ö. 1.1
			1 10 100 1000
			Rat IgG2c Concentration (pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

#### **Background**

There are four IgG subclasses (IgG1, 2, 3 and 4) in Humans, named in order of their abundance in serum (IgG1 being the most abundant). IgG2 is the only IgG subclass which passes through the placenta at a level generally lower than that found in the mother. A deficiency of IgG2 indicates a poor antibody response to bacterial polysaccharides and can lead to increased susceptibility to infections caused by encapsulated bacteria.

Tel: 400-999-2100 Web: <a href="www.elabscience.cn">www.elabscience.cn</a> Email: <a href="techsupport@elabscience.cn">techsupport@elabscience.cn</a>