

## Recombinant Human DYRK3/REDK Protein (His & GST Tag)

**Catalog Number:** PKSH030392

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	Baculovirus-Insect Cells-derived Human DYRK3/REDK protein Met 1-Ser 588, with an N-terminal His & GST
<b>Calculated MW</b>	93.5 kDa
<b>Observed MW</b>	80 kDa
<b>Accession</b>	O43781-1
<b>Bio-activity</b>	The specific activity was determined to be 22 nmol/min/mg using synthetic DYRKtide peptide (RRRFRPASPLRGPPK) as substrate.

### Properties

<b>Purity</b>	> 85 % as determined by reducing SDS-PAGE.
<b>Concentration</b>	Subject to label value.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
<b>Shipping</b>	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
<b>Formulation</b>	Supplied as sterile solution of 20mM Tris, 500mM NaCl, 0.5mM PMSF, 10% glycerol, pH 8.0

### Data



> 85 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Dual specificity tyrosine-phosphorylation-regulated kinase 3, also known as Regulatory erythroid kinase, REDK and DYRK3, is a nucleus protein which belongs to the protein kinase superfamily, CMGC Ser/Thr protein kinase family and MNB/DYRK subfamily. DYRKs are an emerging family of dual-specificity kinases that play key roles in cell proliferation, survival, and development. DYRK3 contains one protein kinase domain. Isoform 1 and isoform 2 of DYRK3 are highly expressed in testis and in hematopoietic tissue such as fetal liver, and bone marrow. Isoform 2 of DYRK3 is the predominant form in testis. Isoform 1 of DYRK3 is the predominant form in fetal liver and bone marrow. Isoform 1 and isoform 2 are present at low levels in heart, pancreas, lymph node, and thymus. DYRK3 is a negative regulator of EPO-dependent erythropoiesis. It may place an upper limit on red cell production during stress erythropoiesis. DYRK3 inhibits cell death due to cytokine withdrawal in hematopoietic progenitor cells. It may also act by regulating CREB/CRE signaling. DYRK3 proved to effectively inhibit NFAT (nuclear factor of activated T cells) transcriptional response pathways and to co-immunoprecipitate with NFATc3. DYRK3 attenuates (and possibly apportions) red cell production selectively during anemia.