

A Reliable Research Partner in Life Science and Medicine

## Recombinant Human DCN Protein (His Tag)

Catalog Number: PDEH100815

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

**Species** Human

Source E.coli-derived Human DCN protein Asp31-Lys359, with an N-terminal His

**Calculated MW** 36.1 kDa Observed MW 38 kDa P07585 Accession

Not validated for activity **Bio-activity** 

#### **Properties**

> 95% as determined by reducing SDS-PAGE. **Purity** 

**Endotoxin** < 10 EU/mg of the protein as determined by the LAL method

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -Storage

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


Shipping This product is provided as lyophilized powder which is shipped with ice packs. **Formulation** 

Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5%

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

#### Data



SDS-PAGE analysis of Human DCN proteins, 2 µg/lane of Recombinant Human DCN proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 38 kDa.

### **Background**

For Research Use Only

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

# Elabscience®

#### Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Decorin is a ubiquitous small cellular or pericellular matrix proteoglycan and is closely related in structure to biglycan protein. It belongs to the small leucine-rich proteoglycan (SLRP) family and consists of a core protein and a covalently linked glycosaminoglycan chain which is either chondroitin sulfate (CS) or dermatan sulfate (DS). As a component of connective tissue, decorin interacts with several extracellular matrix components, such as type I collagen and fibronectin, and plays a role in matrix assembly. Decorin resides in the tumor microenvironment and affects the biology of various types of cancer by downregulating the activity of several receptors involved in cell growth and surviva I. Decorin binds to and modulates the signaling of the epidermal growth factor receptor and other members of the ErbB family of receptor tyrosine kinases. It exerts its antitumor activity by a dual mechanism: via inhibition of these key receptors through their physical downregulation coupled with attenuation of their signaling, and by binding to and sequestering TGFbeta. Decorin also modulates the insulin-like growth factor receptor and the low-density lipoprotein receptor-related protein 1, which indirectly affects the TGFbeta receptor pathway. Decorin plays significant roles in tissue development and assembly, as well as playing both direct and indirect signaling roles.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com