

Recombinant Mouse SCF Protein(His Tag)

Catalog Number: PDMM100227

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	Mammalian-derived Mouse SCF protein Lys26-Ala190, with an C-terminal His
Mol_Mass	18 kDa
Accession	P20826
Bio-activity	Not validated for activity

Properties

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis

Data



SDS-PAGE analysis of Mouse SCF proteins, 2µg/lane of Recombinant Mouse SCF proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 18-38 kDa

Background

For Research Use Only

Tel:400-999-2100

Web:www.elabscience.cn

Email:techsupport@elabscience.cn

Rev. V1.1

Similar to Kit ligand precursor (C-kit ligand), also known as Stem cell factor (SCF), Mast cell growth factor (MGF), or Hematopoietic growth factor KL. SCF/C-kit ligand is the ligand of the tyrosine-kinase receptor encoded by the KIT locus. This ligand is a pleiotropic factor that acts in utero in germ cell and neural cell development, and hematopoiesis, all believed to reflect a role in cell migration. In adults, it functions pleiotropically, while mostly noted for its continued requirement in hematopoiesis. SCF/C-kit ligand stimulates the proliferation of mast cells. This protein can augment the proliferation of both myeloid and lymphoid hematopoietic progenitors in bone marrow culture. It may act synergistically with other cytokines, probably interleukins. SCF/C-kit ligand is the ligand for the tyrosine kinase receptor c-kit, which is expressed on both primitive and mature hematopoietic progenitor cells. In vitro, SCF/C-kit ligand synergizes with other growth factors, such as granulocyte colony-stimulating factor (G-CSF), granulocyte-macrophage colony-stimulating factor, and interleukin-3 to stimulate the proliferation and differentiation of cells of the lymphoid, myeloid, erythroid, and megakaryocytic lineages. In vivo, SCF/C-kit also synergizes with other growth factors and has been shown to enhance the mobilization of peripheral blood progenitor cells in combination with G-CSF. In phase I/II clinical studies administration of the combination of SCF and G-CSF resulted in a two- to threefold increase in cells that express the CD34 antigen compared with G-CSF alone.