Recombinant Human CDKN1B Protein (His Tag)

Catalog Number: PKSH032314

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | | | | |
|----------------|--|--|--|--|
| Species | Human | | | |
| Source | E.coli-derived Human CDKN1B protein Met 1-Thr198, with an N-terminal His | | | |
| Calculated MW | 24.2 kDa | | | |
| Observed MW | 30 kDa | | | |
| Accession | P46527 | | | |
| Bio-activity | Not validated for activity | | | |
| Properties | | | | |
| Purity | > 95 % as determined by reducing SDS-PAGE. | | | |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. | | | |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 | | | |
| | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of | | | |
| | reconstituted samples are stable at $< -20^{\circ}C$ for 3 months. | | | |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. | | | |
| Formulation | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. | | | |
| | Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants | | | |
| | before lyophilization. | | | |
| | Please refer to the specific buffer information in the printed manual. | | | |
| Reconstitution | Please refer to the printed manual for detailed information. | | | |

Data

| kDa 120 90 60 40 | MK | R | |
|------------------------------|----|---|--|
| 30 | - | | |
| 20 | - | | |
| 14 | _ | | |

> 95 % as determined by reducing SDS-PAGE.

Background

Cyclin-Dependent Kinase Inhibitor 1B (CDKN1B) is a Kinesin-related motor protein necessary for mitotic spindle assembly and chromosome segregation. CDKN1B is expressed in all tissues with highest levels observed in skeletal muscle. CDKN1B is a potent inhibitor of Cyclin E- and Cyclin A-CDK2 complexes. CDKN1B forms a complex with Cyclin Type D-CDK4 complexes and is involved in the assembly, stability, and modulation of CCND1-CDK4 complex activation. In addition, CDKN1B acts as an inhibitor or an activator of Cyclin Type D-CDK4 complexes depending on its phosphorylation state and stoichometry.

Web:www.elabscience.com