

## APC Anti-Rat CD3 Antibody[G4.18]

Catalog Number: E-AB-F1228E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Reactivity	Rat
Host	Mouse
Isotype	Mouse IgG3, κ
Clone No.	G4.18
Isotype Control	APC Mouse IgG3, κ Isotype Control[A112-3] [Product E-AB-F09752E]
Conjugation	APC
Conjugation Information	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.


### Applications

### Recommended usage

#### FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### Data



Rat splenocytes are stained with APC Anti-Rat CD3 Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

### Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

### Antigen Information

Alternate Names	T3and ζ chains;CD3;CD3 Complex;T-cell surface glycoprotein CD3 δ
Uniprot ID	P19377;Q64159;D4A5M2
Gene ID	25710,300678,315609,25300

### For Research Use Only

## Background

CD3 is a complex composed of  $\delta$ ,  $\gamma$ ,  $\epsilon$ , and  $\zeta$  chains. They are 20-25 kD members of the immunoglobulin superfamily and associated with the T cell receptor (TCR). CD3 is expressed on thymocytes, peripheral T cells, some NK-T cells, and dendritic epidermal T cells. CD3 is involved in antigen recognition, signal transduction, and T cell activation