



A Reliable Research Partner in Life Science and Medicine

# **Human COMP Antibody Pair Set**

Catalog No. E-KAB-0261 Applications ELISA

**Synonyms** EDM1, EPD1, MED, PSACH, THBS5, TSP-5

## **Kit components & Storage**

Title	Specifications	Storage
Human COMP Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Human COMP Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### **Product Information**

Items		Characteristic (E-KAB-0261)	
		Human COMP Capture Antibody	Human COMP Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Human COMP protein	Recombinant Human COMP protein
Information	Swissprot	P49747	
Product details	Reactivity	Human	Human
	Host	Mouse	Rabbit
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
-			7.4
	Purify	Protein A	Protein A & Antigen Affinity
	Specificity	Detects Human COMP in ELISAs.	·

For Research Use Only

Tel: 400-999-2100 Web: <a href="www.elabscience.cn">www.elabscience.cn</a> Email: <a href="mailto:techsupport@elabscience.cn">techsupport@elabscience.cn</a>





A Reliable Research Partner in Life Science and Medicine

## **Applications**

Human COMP Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Human COMP Capture Antibody	
Capture			Aig
ELISA Detection	1:1000-1:10000	Human COMP Detection Antibody (Biotin)	0.01 0.01 0.01 1 10 100 1000 Human COMP concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

#### **Background**

The protein encoded by this gene is a noncollagenous extracellular matrix (ECM) protein. It consists of five identical glycoprotein subunits, each with EGF-like and calcium-binding (thrombospondin-like) domains. Oligomerization results from formation of a five-stranded coiled coil and disulfides. Binding to other ECM proteins such as collagen appears to depend on divalent cations. Mutations can cause the osteochondrodysplasias pseudochondroplasia (PSACH) and multiple epiphyseal dysplasia (MED).

Tel: 400-999-2100 Web: <a href="www.elabscience.cn">www.elabscience.cn</a> Email: <a href="mailto:techsupport@elabscience.cn">techsupport@elabscience.cn</a>