

Recombinant CD180/RP105 Monoclonal Antibody

catalog number: **AN300524P**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Mouse
Immunogen	Recombinant Mouse CD180/RP105 Protein
Host	Rabbit
Isotype	IgG
Clone	8A3
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS


Applications Recommended Dilution

IHC-P	1:50-1:200
--------------	------------

Data



Immunohistochemistry of paraffin-embedded mouse thymus using CD180/RP105 Monoclonal Antibody at dilution of 1:100.



Immunohistochemistry of paraffin-embedded mouse spleen using CD180/RP105 Monoclonal Antibody at dilution of 1:100.

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD180, also known as RP105, is a B-cell surface molecule belonging to the family of pathogen receptors, Toll-like receptors (TLR). CD180 has an extracellular leucine-rich repeats and a short cytoplasmic tail. CD180/RP105 interacts with an extracellular molecule named MD1 and then together form the cell surface receptor complex RP105/MD1 which induces B-cell activation in humans and mice, leading to proliferation and up-regulation of a costimulatory molecule, B7.2/CD86. CD180/RP105 also has a role in LPS response because B cells lacking RP105 show hyporesponsiveness to LPS.

For Research Use Only